

CONTENTS

| | |
|--|-----|
| Conformational analysis of β -D-fructofuranosyl-(2 \rightarrow 6)- β -D-glucopyranoside by molecular mechanics (MM2) calculations A.L. Waterhouse (Davis, CA, USA), K. Horváth and J. Liu (New Orleans, LA, USA) . . . | 1 |
| Observation by solid-state ^{13}C CP MAS NMR spectroscopy of the transformations of wheat starch associated with the making and staling of bread K.R. Morgan, R.H. Furneaux and R.A. Stanley (Lower Hutt, New Zealand) | 15 |
| X-ray fibre diffraction studies of chitosan and chitosan gels P. Cairns, M.J. Miles, V.J. Morris, M.J. Ridout, G.J. Brownsey (Norwich, United Kingdom) and W.T. Winter (Syracuse, NY, USA) | 23 |
| Analysis of residue types in curdlan sulfate by nuclear magnetic resonance H. Miyano, R. Nakagawa, E.-i. Suzuki (Kawasaki, Japan) and T. Uryu (Tokyo, Japan) . . . | 29 |
| On the characterization of the reaction of organotin compounds with D-glucuronic acid H.E. Guard, W.M. Coleman III (Arlington, VA, USA) and M.M. Ross (Washington, DC, USA) | 41 |
| A ^1H NMR database computer program for the analysis of the primary structure of complex carbohydrates J.A. van Kuik, K. Hård and J.F.G. Vliegthart (Utrecht, Netherlands) | 53 |
| A ^1H and ^{13}C NMR study of oligosaccharides from human milk. Application of the computer program CASPER K. Hermansson, P.-E. Jansson, L. Kenne, G. Widmalm and F. Lindh (Stockholm, Sweden). | 69 |
| Fmoc-protected, glycosylated asparagines potentially useful as reagents in the solid-phase synthesis of N-glycopeptides L. Urge, L. Otvos, Jr., E. Lang, K. Wroblewski (Philadelphia, PA, USA), I. Laczko (Szeged, Hungary) and M. Hollosi (Budapest, Hungary) | 83 |
| Stereoselective total synthesis of the glycosyl phosphatidylinositol (GPI) anchor of <i>Trypanosoma brucei</i> C. Murakata (Saitama, Japan) and T. Ogawa (Tokyo, Japan) | 95 |
| Large-scale synthesis of D-mannose 6-phosphate and other hexose 6-phosphates M. Meldal, M.K. Christensen and K. Bock (Valby, Denmark) | 115 |
| Synthesis of a cycloheptaose consisting of (1 \rightarrow 4)-linked 7-amino-6,7-dideoxy- α -D-glucopyranosyl units: A new analog of cyclomaltoheptaose H.H. Baer, Y. Shen (Ottawa, ON, Canada), F. Santoyo González, A. Vargas Berenguel and J. Isac García (Granada, Spain) | 129 |

| | |
|--|-----|
| Synthesis and collagenase inhibition of new glycosides of aranciamycinone: the aglycon of the naturally occurring antibiotic aranciamycin M. Bols (Lyngby, Denmark), L. Binderup, J. Hansen and P. Rasmussen (Ballerup, Denmark) | 141 |
| Synthesis of 4-C-methyl analogues of glucosylceramide M. Plewe (Konstanz, Germany), K. Sandhoff (Bonn, Germany) and R.R. Schmidt (Konstanz, Germany) | 151 |
| Synthesis, crystalline structure, conformational analysis, and azidolysis of methyl 2,3-anhydro- α -D-manno- and -allo-pyranoside <i>p</i> -bromobenzyl ethers X. Wu, F. Kong, D. Lu and G. Li (Beijing, China) | 163 |
| Sulfated sialyl-oligosaccharides derived from tracheobronchial mucous glycoproteins of a patient suffering from cystic fibrosis. T.P. Mawhinney, D.C. Landrum, D.A. Gayer and G.J. Barbero (Columbia, MO, USA) . . | 179 |
| The capsular antigen of <i>Escherichia coli</i> serotype O8:K102:H ⁻ A.H. De Bruin, H. Parolis and L.A.S. Parolis (Grahamstown, South Africa) | 199 |
| The structure of the capsular polysaccharide of <i>Escherichia coli</i> O9:K35:H ⁻ P.L. Hackland and H. Parolis (Grahamstown, South Africa) | 211 |
| The structure of the fructan sinistrin from <i>Urginea maritima</i> T. Spies, W. Praznik, A. Hofinger, F. Altmann (Wien, Austria), E. Nitsch and R. Wutka (Linz, Austria) | 221 |
| Fructo-oligosaccharides from the stems of <i>Triticum aestivum</i> W. Praznik, T. Spies and A. Hofinger (Wien, Austria) | 231 |
| Quantitative analysis of carboxymethyl chitin adsorbed on a liposome surface T. Nishiya and D. Robibo (Montreal, PQ, Canada) | 239 |
| Development of a water-soluble, sulfated (1 \rightarrow 3)- β -D-glucan biological response modifier derived from <i>Saccharomyces cerevisiae</i> D.L. Williams (Johnson City, TN, USA), H.A. Pretus, R.B. McNamee, E.L. Jones, H.E. Ensley (New Orleans, LA, USA) and I.W. Browder (Johnson City, TN, USA) | 247 |
| ¹ H-NMR-Untersuchungen zur Spaltspezifität von α - und β -D-Glucosidasen gegenüber Disacchariden. II R. Wajda, M. König, H. Ludwig, S. Mronga und H. Friebolin (Heidelberg, Deutschland) . | 259 |
| Mechanism of action of D-galacturonan digalacturonohydrolase of <i>Selenomonas ruminantium</i> on oligogalactosiduronic acids K. Heinrichová, M. Dzúrová and L. Rexová-Benková (Bratislava, Czechoslovakia) | 269 |
| <i>Notes</i> | |
| Nouvelle voie d'accès au D-ribose et au D-lyxose B. Lacourt-Gadras, M. Grignon-Dubois et B. Rezzonico (Talence, France) | 281 |
| Synthesis of <i>N</i> -acetyl-lactosamine via ozonolysis of a nitro derivative E. Lattová and L. Petruš (Bratislava, Czechoslovakia) | 289 |

| | |
|---|-----|
| A simple method for avoiding alkylthio group migration during the synthesis of thioglycoside 2,3-orthoesters. An improved synthesis of partially acylated 1-thio- α -L-rhamnopyranosides V. Pozsgay (Bethesda, MD, USA) | 295 |
| Isolation and identification of <i>O</i> - β -D-fructofuranosyl-(2 \rightarrow 1)- <i>O</i> - β -D-fructofuranosyl-(2 \rightarrow 1)-D-fructose, a product of the enzymic hydrolysis of the inulin from <i>Cichorium intybus</i> A. De Bruyn, A.P. Alvarez, P. Sandra (Gent, Belgium) and L. De Leenheer (Tienen, Belgium) | 303 |
| Studies on anticoagulant-active arabinan sulfates from the green alga, <i>Codium latum</i> T. Uehara, M. Takeshita and M. Maeda (Saitama, Japan) | 309 |
| <i>Preliminary communication</i> | |
| One- and two-dimensional ^1H NMR investigations of the inclusion of the anti-cancer drug mitoxantrone in cyclomaltooligosaccharides N. Birlirakis and B. Perly (Gif sur Yvette, France) | C1 |
| A new method for <i>N</i> -alkyl and <i>N,N</i> -dialkyl-D-glucosamines J.M. Vega-Pérez, J.L. Espartero and F. Alcudia (Sevilla, Spain) | C5 |
| A study of possible sulfate loss during the chemical release of sulfated oligosaccharides from glycoproteins K.R. King, J.M. Williams (Swansea, United Kingdom), J.R. Clamp and A.P. Corfield (Bristol, United Kingdom) | C9 |
| A facile, systematic synthesis of ganglio-series gangliosides: Total synthesis of gangliosides GM ₁ and GD _{1a} A. Hasegawa, T. Nagahama and M. Kiso (Gifu, Japan) | C13 |
| Structure of the O-specific polysaccharide of <i>Proteus penneri</i> 62 containing 2-acetamido-3- <i>O</i> -[(<i>S</i>)-1-carboxyethyl]-2-deoxy-D-glucose (<i>N</i> -acetylisomuramic acid) Y.A. Knirel, N.A. Paramonov, E.V. Vinogradov, A.S. Shashkov, N.K. Kochetkov (Moscow, Russian Federation), Z. Sidorczyk and A. Swierzko (Lodz, Poland) | C19 |
| <i>Announcements</i> | C25 |
| <i>Author index</i> | C27 |
| <i>Subject index</i> | C29 |
| <i>Contents</i> | C33 |

